



ABS051

EVALUATION OF MODIFIED BORG DYSPNOEA SCORE AND 6-MINUTE WALK TEST IN PULMONARY ARTERIAL HYPERTENSION PATIENTS RECEIVING PHOSPHODIESTERASE INHIBITORS

Teena Nazeem^{*1},
Sabin Thomas²,
Dr. Govinda Ajmera³

¹Assistant Professor,
Krupanidhi College of
Pharmacy, Bangalore

²Assistant Professor,
College of Pharmacy,
University of Nizwa,
Sultanate of Oman

³Clinical Pharmacist,
Clinical Pharmacy
Department, CHL Apollo,
Indore.

Introduction: Pulmonary Arterial Hypertension (PAH) is a progressive condition that results in right heart failure and death¹. Patients with PAH who do not receive disease targeted therapy have a poor quality of life and high mortality rates with a median survival of less than three years from diagnosis^{2,3}. In the last few years, oral Sildenafil Citrate (a specific phosphodiesterase-5 inhibitor which is widely used in the treatment of erectile dysfunction), has shown promising results as a novel oral monotherapy in the treatment of PAH^{4,5}.

Methodology: A prospective open label randomized study was conducted over a period of 8 months in hospitalized patients who visited the in-patient block in a tertiary care hospital in Nilgris. Simple Randomization technique was adopted to allocate the patients into three study groups. At the baseline Visit, Modified Borg Dyspnea Scale (MBDS) and 6 MWT (6 Minute Walk Test) were evaluated to confirm study criteria and were allotted into 3 doses based on their severity upon Right Ventricular Systolic Pressure. The same parameters were assessed on each visit for all the patients throughout the study period.

Results and Discussion: In our study, Sildenafil Citrate showed an improvement in MBDS and 6 MWT after 8 months of therapy which were assessed using Kruskal-Wallis non-parametric test. A remarkable improvement in 6 MWT from 171.88 $\pm$ 52.4 metres to 337.12 $\pm$ 71.84 metres was observed. There was an improvement in dyspnea by Borg scale in all study groups. Cor-pulmonale patients were more benefited ($p < 0.001$) whereas patients with Atrial Septal Defect (ASD) and Rheumatic Heart Disease (RHD) also improved significantly ($p < 0.05$).

Conclusion: Sildenafil has been reported to be well tolerated among the patient population. The patients who were slightly affected at baseline with moderate to severe dyspnoea became slight grade which

Keywords: PAH, MBDS, 6MWT