

REVIEW ON NIOSOMES - A NOVEL APPROACH FOR DRUG TARGETING

Preethi Sudheer and Kar Kaushik

Krupanidhi College of Pharmacy, #12/1, Chikkabellandur, Carmelaram Post, Bangalore - 560 035, Karnataka, India. Mob. No: +919449822912.

Received on: 19.02.2015

Revised: 11.03.2015

Accepted: 17.03.2015

ABSTRACT

Purpose: This is a general review on niosome as drug carrier, which improves the bioavailability of drugs, increases penetration through skin, releases drug in a controlled or sustained manner, and used for targeting of drugs to specific sites in the body.

Approach: Niosomal drug delivery system can be considered as an emerging novel drug delivery system, which consists of microscopic non-ionic vesicles composed of non-ionic surfactants. These are biodegradable, relatively nontoxic, more stable and inexpensive, and an alternative to liposomes. Methods of preparation, characterisation and application of niosomes have been reviewed.

Findings: Niosome has potential to reduce the side effects of drugs and increase therapeutic effectiveness in various diseases. It can also be used as a carrier to deliver drugs topically.

Conclusion: This review presents an overview of the types of niosomes, techniques of preparation of niosome, characterisation and their applications.

Keywords: Niosome; non-ionic surfactant; drug entrapment; novel drug delivery system.

INTRODUCTION

Since past many years, research is going on for finding unique and better alternatives of drug delivery system and this will continue till the drug delivery system have no side effects with optimum therapeutic action. Conventional dosage forms are still in use because of high patient compliance, though they are having side effects. Recently, vesicular systems which have lot of advantages over other drug delivery systems are being extensively explored. Vesicular system includes niosome, liposome, transferosomes. Niosome or non-ionic surfactant vesicles are those which act as drug carrier to deliver drug to the site of action. Niosomes are vesicular carriers of non-ionic surfactants with size range between 10 and 1000 nm, wherein aqueous phase is enclosed in a highly ordered bilayer of non-ionic surfactants with or without cholesterol and dicetyl phosphate. Niosomes can be used for entrapping both hydrophilic and hydrophobic drugs (Fig 1).




Fig. 1: Structure of niosomes (reproduced from Vedha Hari BN, et al. Novel technologies: A weapon against tuberculosis. Ind J col. 2010;6: 338-344.)

Niosomes are novel drug delivery system, preferred over other vesicular system because of its low toxicity, non ionic nature, bio-degradable, better availability of drug at site, and good intrinsic skin penetration. Encapsulating the drug in the niosome, can protect the drug from acidic and enzymatic degradation after administration.¹

Many attempts have been made in the research of niosomes to encapsulate various NSAIDs for better bioavailability, and better permeation through the skin.

Novel drug delivery system has limitations such as high cost, which makes it difficult for production and have problem with dose adjustments. So this problem can be overcome by using vesicular drug delivery system like niosomes, which may prolong the existence of the drug in the systemic circulation, increase permeation through skin and reduce toxicity.²

Advantages of Niosomes

Niosomes can act as a carrier for drugs which have a wide range of solubility as it has both hydrophilic and hydrophobic part.

Surfactants used in niosome are bio-degradable, biocompatible, and nonimmunogenic.

Niosomes of desired property can be prepared by altering vesicle characteristics such as vesicle composition, size, lamellarity, tapped volume, surface charge.

Niosomes can also be used as a depot formulation.

As niosomal suspension has hydrophilic tail, therefore it can be considered to provide better patient compliance than oil based formulations.

*Correspondence : email : preetisudheer@gmail.com

Review on Niosomes

Niosomes not only have good stability but also have osmotic property which makes it superior to oil based formulations and also increases the entrapment efficiency.

Niosomes are considered to be versatile as it can be given by oral, parenteral as well as topical routes.

Niosomes are proposed to penetrate through the cornea i.e. it can be used for ocular drug delivery.^{3,4}

Disadvantages of Niosomes

Niosomal aqueous suspensions owe limited shelf life due to fusion, aggregation, leaking of entrapped drugs, and hydrolysis of encapsulated drugs.

The techniques involved in the niosomal formulation such as extrusion, sonication are time consuming and requires specialized equipments for processing.⁵

Various Types of Niosome

Niosomes can be divided into three groups based on the vesicle size. These are –

Small unilamellar vesicles= 0.025- 0.05 μm .

Multilamellar vesicles = >0.05 μm .

Large unilamellar vesicles= >0.10 μm .

METHODS OF PREPARATION

Niosomes are prepared by various methods and they are:

Preparation of Small Unilamellar Vesicles

Sonication: This method involves addition of surfactant/cholesterol in organic solvent and drug in aqueous solution. This aqueous solution of drug is mixed with surfactant solution and further homogenized for 3 minutes at a temperature of 60°C.⁶

Micro fluidization: Submerged jet principle is adopted for niosome formulation. Fluidised stream of drug and surfactant is allowed to interact at ultra high speed velocities through micro channel within the interaction chamber. The impingement of thin liquid sheet along a common front is arranged such that the energy utilised by the system remains within the area of niosome formation. The method results in the formation of niosomes of greater uniformity, smaller size and better reproducibility.⁵

Preparation of Multilamellar Vesicles

Hand shaking method (Thin film hydration technique): This method involves formation of thin dried layer in the rota flask. In this method, surfactant and cholesterol is dissolved in organic solvent. Both the solutions are mixed and the mixture is evaporated under reduced pressure at definite temperature to form thin layer on the inner wall of the flask. After evaporation, hydration is carried out with sonication to form niosomes.⁶

Trans-membrane pH gradient drug uptake process: This method utilizes thin film hydration technique. In this method, a definite ratio of cholesterol and surfactant is

Preethi Sudheer and Kaushik Kar

dissolved in organic solvents like ether or chloroform and evaporated under reduced pressure to form thin dried film. The resultant film is hydrated with 300 mM citric acid (pH 4.0) by vortex mixing. The vesicles formed are allowed for freezing and thawing several times and sonicated to get niosomes. Aqueous solution of drug is added to the above niosomal suspension and is vortexed. The pH of niosomal suspension is adjusted to 7.0-7.2 with 1 M disodium phosphate, heated to 60°C for 10 minutes to form the desired multilamellar vesicles.⁷

Preparation of Large Unilamellar Vesicles

Reverse phase evaporation technique (REV): In this method, a definite ratio of cholesterol and surfactant is dissolved in organic solvents like ether or chloroform and evaporated under reduced pressure to form thin dried film. The resultant film is hydrated with 300 mM citric acid (pH 4.0) by vortex mixing. The gel formed is further sonicated after addition of small amount of phosphate buffer solution and the organic phase present is removed at 40°C which forms niosomes having high viscosity. This viscous niosome is diluted with phosphate buffer saline which is maintained at 60°C for 10-15 minutes.⁸

Ether injection method: This method is used in the preparation of large unilamellar vesicles, where definite ratio of cholesterol and surfactant is dissolved in organic solvent. To the above mixture, organic solution of drug is added and injected to the aqueous solution positioned at magnetic stirrer maintained at 60°C with continuous stirring forming niosomal suspension.⁹

Miscellaneous

Multiple membrane extrusion method: In this method surfactant and cholesterol are dissolved in organic solvent i.e. chloroform. Then dicetyl phosphate is added to the organic solution of surfactant/cholesterol, and the resultant solution is evaporated under reduced pressure to form thin film. The film formed is rehydrated with aqueous solution of drug. The resultant suspension is extruded through polycarbonate membrane, placed in a series for up to eight passages. Size of niosomal formulation can be controlled by this method.¹⁰

Emulsion method: An aqueous solution of drug is prepared separately and this solution is added to the solution of surfactant and cholesterol dissolved in organic solvent, forming oil in water emulsion. The organic solvent is then evaporated leaving niosomes dispersed in the aqueous phase.¹¹

Lipid Injection Method: This method uses lipids along with surfactants. Lipids and surfactants are melt and the mixture is injected into aqueous solution of drug which is heated. Another method involves addition of drug in lipids and melting. The molten solution is injected into an aqueous solution of surfactant which is heated to form niosomes.¹²

The bubble method: This is a one step technique for the preparation of niosome without the use of organic solvent. The bubbling unit has round-bottomed flask with

Review on Niosomes

three necks positioned in water bath for controlling the temperature. In the first neck, water cooled reflux is placed, thermometer placed in the second and nitrogen supply is provided through the third neck. Dispersion of niosomes is prepared by adding cholesterol and surfactant in phosphate buffer pH 7.4 which is maintained at 70°C and using high shear homogenizer the dispersion formed is mixed for 15 seconds, then it is bubbled using nitrogen gas at 70°C.¹³

SEPARATION OF UNENTRAPPED DRUG

The untrapped drug can be separated by various techniques:

Dialysis: Free drug is separated from niosomal dispersion by using a dialysis bag in phosphate buffer as media.

Gel Filtration: The untrapped drug in niosomal dispersion is removed by using gel filtration by passing through Sephadex G-50 column and eluted with phosphate buffer or normal saline.

Centrifugation: Here the separation of the free drug is achieved by centrifugation of the niosomal suspension for 30 minutes. The supernatant solution is decanted, and the precipitate is washed and resuspended in phosphate buffer to obtain a niosomal suspension free from untrapped drug.¹⁴

CHARACTERISATION OF NIOSOMES

Vesicle diameter: The shape of niosomes is spherical in nature and the size ranges from 20 nm to 50 nm. Vesicular form and size distribution can be determined by light microscopy. Another method used for determination of diameter and size of niosome is freeze fracture electron microscopy.¹⁵

Vesicle charge: Vesicle charge is determined by measuring zeta potential. Charged niosomes are more stable than uncharged niosomal dispersion. Hence, during the preparation, charged vehicles like dicetyl phosphate is added to surfactant/cholesterol mixture.¹⁶

Bilayer formation & number of lamellae: Bilayer formation of niosomes is characterized by X-cross formation. NMR spectroscopy, electron microscopy and small angle X-ray scattering is used to characterize number of lamellae.^{17,18}

Membrane rigidity and homogeneity: Rigidity of membrane affects bio-degradation and bio-distribution of niosomes. Determination of rigidity of niosomal suspension is done by fluorescence probe as a function of temperature. P-NMR, differential scanning calorimetry (DSC), Fourier transform-infra red spectroscopy (FTIR) and fluorescence resonance energy transfer (FRET) are used to determine the membrane homogeneity.¹⁸

Encapsulation efficiency: It can be done by separating the untrapped drug in the niosomal dispersion by using various methods like centrifugation, gel filtration and dialysis. Other method of separation includes complete disruption of vesicle using 50% n-propanol or

Preethi Sudheer and Kaushik Kar

0.1% Triton X-100. Then the required disrupted vesicle is analysed for the drug content and hence entrapment efficiency.¹⁹

Entrapment efficiency (EF) = (Amount entrapped/ total amount) x 100

In vitro drug release: *In vitro* drug release of niosomes can be characterized by the following methods:

1. Dialysis
2. Franz diffusion cell

Dialysis: Dialysis is carried out by placing niosomal dispersion in dialysis bag which is pre-washed and pre-soaked and are tied at both the ends. The dialysis bag is suspended into a dissolution media maintained at 37 °C with continuous stirring. Samples are withdrawn at regular time interval and replaced with fresh sample. The withdrawn samples are analyzed for the drug content.²⁰

Franz diffusion cell: This cell assembly consists of donor and receptor compartment which is separated by cellophane membrane or dialysis membrane. Niosomal dispersion is placed at donor compartment and dissolution media i.e. phosphate buffer is placed in receptor compartment maintained at 37°C with continuous stirring by magnetic stirrer. The samples are taken out at suitable time intervals, replaced with fresh media and analyzed for drug content.²¹

In vivo study: The *in vivo* study is carried out in albino wistar male rats weighing 150-200 gm. The animals are divided into 3 groups, each group containing 6 animals. Group I is treated as treatment control, group II with pure drug solution, and group III with niosomal formulation of the drug, (dose to be determined). Blood samples are collected from the rats at the interval of 0.5h, 2h, 4h, 8h, 12h, and 16 h, from the catheter, which is implanted in the femoral artery. The catheter is continuously flushed with physiological saline. The collected blood samples are centrifuged in a cooling centrifuge at 12,000 rpm for 3min and blood plasma is separated. The separated blood plasma is stored in a freezer at -4°C until it is analysed by HPLC or any other precise method.²²

Stability of niosomes: Niosomes are stored at two different conditions, usually 4±1° and 25±2°. The constant particle size and constant concentration of entrapped drug indicates stability of niosomes. The concentration and type of surfactant, cholesterol also affects niosome stability. Light microscope is used for the determination of vesicle size and the number of vesicles per cubic mm.²³

FACTORS GOVERNING NIOSOME FORMATION

Choice of surfactants and additives: Non-ionic surfactants are employed for formation of niosome vesicles. Surfactants having hydrophobic tail may consist of one or two alkyl or perfluoroalkyl groups or, in some cases, a single steroidal group. The ether-type surfactants with mono alkyl chain are more toxic than ester-type surfactants. When stability comes into consideration, ester type surfactant is less stable than ether-type surfactants and it is because, ester linked surfactants get degraded by esterase into triglycerides

Review on Niosomes

and fatty acid, *in vivo*. The surfactants with alkyl chain length from C₁₂ to C₁₈ are suitable for preparation of niosome. Niosome vesicle formation finds suitability in surfactants having HLB value between 4 and 8.^{24,25}

Surfactant and lipid level: Generally surfactant/lipid level which is essential for niosomal formulation is kept between 10-30 mM (1-2.5% w/w). The niosomal dispersion is affected by alteration of surfactant, water ratio during the hydration step. By increasing surfactant/lipid level the total amount of drug encapsulated also increases.²⁶

Composition of membrane: Stabilisation of niosomes can be achieved by adding different additives to the surfactant mixture. One main disadvantage of niosome formulation is the leakage of drug from the vesicles which can be controlled by addition of cholesterol. Cholesterol confers better rigidity to the membrane and therefore leakage of drug is reduced.¹³

Nature of encapsulated drug: Niosomal formulation is influenced by the nature of drug being encapsulated. The physico-chemical properties of the encapsulated drug influence charge and the rigidity of the niosomal bilayer. The interaction of the surfactant head groups lead to entrapment of drug in vesicles and cause increase in charge. The formation of charge creates mutual repulsion of the surfactant bilayer and thus increases vesicle size. The HLB of drug also affects the degree of entrapment.²⁷

Temperature of hydration: The shape and size of niosome is influenced by hydration temperature. Variation in temperature affects surfactants to assemble into vesicles and therefore, affects niosomal vesicle formation. The hydration temperature for niosome formation should be above the gel to liquid phase transition temperature of system.^{23,24}

Cholesterol content: Addition of cholesterol in niosomal formulation improves entrapment efficiency and thereby provides rigidity to the vesicles. It also increases the hydrodynamic diameter of the niosomal vesicles. Cholesterol also increases the chain order of liquid state bilayers and decreases the chain order of gel state bilayers. Gel state bilayers can be converted into liquid state by increasing the concentration of cholesterol.²⁸

Charge: In multilamellar vesicle structure, the interlamellar distance between successive bilayers increases due to presence of charge. This results in greater overall entrapped volume.²⁰

APPLICATIONS

Generally peptide drugs have stability problem so it is difficult to formulate in the form of tablets, parenteral. Therefore, using niosome as a drug carrier can improve the stability of peptide drugs.^{29,30}

Nowadays due to various disadvantages of oral drug delivery system, research is going on for transdermal drug delivery and this has achieved good response. Niosome as a drug

Preethi Sudheer and Kaushik Kar

carrier has good penetration capacity. Therefore, niosome can be used as transdermal drug delivery for various drugs.³¹

Niosomes can be used as a suitable drug delivery system for the administration of anticancer drugs such as 5-FU cancer therapy. It is also used for increasing the efficacy of drug by incorporating it into niosome.³²

Niosomal suspension shows a visible spectrum which is superimposable onto that of free haemoglobin; therefore it can be used as a carrier for haemoglobin.³³

Niosomal system can encapsulate drugs which have low therapeutic index and low water solubility and this can be maintained in the circulation showing sustained release action.³³

The treatment of leishmaniasis is mostly achieved by antimony derivatives. It has side-effects at higher concentrations which may cause liver, cardiac and kidney damage. These side effects at higher concentration can be overcome by niosomes as a drug carrier.³³

Niosomal system also finds use in diagnostic agents. It can also act as a carrier for radiopharmaceuticals.³⁴

Niosomal formulation of diclofenac sodium prepared with 70% cholesterol showed greater anti-inflammatory effect as compared to the free drug. Similarly, nimesulide and flurbiprofen showed greater activity than the free drug.³⁵

RECENT DEVELOPMENTS

Various studies have been carried out in the field of niosomes. Attempts were made to improve the oral bioavailability of Cefdinir by incorporating into niosomes and study revealed that formulated niosome showed higher release and improved permeability across animal intestinal membrane than plain drug formulation and marketed formulation.³⁶

Research carried out to increase the solubility of BCS class II drug valproic acid by incorporating into a niosomal gel concluded that niosomal entrapped gel had better *in vitro* release across the animal nasal membrane and hence it can be used as an efficient carrier for valproic acid.³⁷

Oxcarbazepine niosomes were formulated and evaluated to explore the advantage over the conventional dosage forms. Pharmacokinetic studies have shown that oxcarbazepine niosomes have increased elimination half life and the area under the curve also higher when compared with pure oxcarbazepine, which justify their potential in strengthening the efficacy and safety profile of the drug.²²

CONCLUSION

The field of vesicular drug delivery system is still in its infancy and increasing gradually during the past few decades. It is expected that this trend will continue to increase further. Niosome is a promising vesicular

Review on Niosomes

delivery system compared to liposomes because of its low cost, stability and the ability to encapsulate different type of drugs. Researches have shown that it is useful in the delivery of anti-cancer, anti epileptic and anti-inflammatory agents. It also targets certain areas of the mammalian anatomy and hence exploited as diagnostic imaging agent. It has expanded to various areas like vaccine delivery system, tumour targeting agents, ophthalmic, and transdermal delivery systems. Hence, further research has to be carried out to extract out the full potential of niosomes.

ACKNOWLEDGMENTS

The authors thank the Management, the Principal and Head of the Department of Pharmaceutics, Krupanidhi College of Pharmacy, Bangalore for their encouragement.

CONFLICT OF INTEREST

The authors report that there is no conflict of interest.

REFERENCES

1. Yie W Chien. Novel Drug Delivery System, 2nded. New York: Marcel Dekker Inc; 1992. p. 1-42.
2. Tsuchihashi M, Harashima H, Kiwada H. Development of pharmacokinetic pharmacodynamic (PK/PD)-simulation system for doxorubicin in long circulating leptosomes in mice using peritoneal. J control Rel. 1999;61:9-19.
3. Biju SS, Telegaonar S, Mishra PR, Khar RK. Vesicular system: an overview. Ind J Pharm Sci. 2006;68:141-53.
4. Vedhahari BN, Chitra KP, Bhimavarapu R, Karunakaran P, Muthukrishnan N, Rani B, et.al Novel technologies: A weapon against tuberculosis. Ind J Pharmacol. 2010;6:338-344.
5. Verma S, Singh SK, Navneet S, Mathur P, Valecha V. Nanoparticle vesicular systems: a versatile tool for drug delivery. J Chem Pharm Res. 2010; 2(2):496-509.
6. Baillie AJ, Coombs GH, Dolan TF, Laurie J. Non-ionic surfactant vesicles, niosomes as delivery system for the anti-leishmanial drug, sodium stibogluconate. J Pharmacol. 1986; 38:502-5.
7. Mayer LD, Bally MB, Hope MJ, Cullis PR. Uptake of antineoplastic agents into large unilamellar vesicles in response to a membrane potential. Biochem Biophys Acta. 1985;816: 294-302.
8. Raja RA, Chandrashekhar G, Pillai GK, Udupa N. Antiinflammatory activity of niosome encapsulated diclofenac sodium with tween -85 in arthritic rats. Ind J Pharmacol. 1994;26:46-48.
9. Rogerson A, Cummings J, Willmott N, Florence AT. The distribution of doxorubicin in mice following administration in niosomes. J Pharm Pharmacol. 1988;40:337-42.

Preethi Sudheer and Kaushik Kar

10. Jayaraman SC, Ramachandran C, Weiner N. Topical delivery of erythromycin from various formulations: an *in vivo* hairless mouse study. J Pharm Sci. 1996;85:1082-84.
11. Hao Y, Zhao F, Li N, Yang Y, Li K. Studies on a high encapsulation of colchicines by a niosome system. Int J Pharm. 2002;244:73-80.
12. Karim KM, Mandal AS, Biswas N, Guha A, Chatterjee S, Behera M. Niosome: a future of targeted drug delivery systems. J Adv Pharm Tech Res. 2011;62:122.
13. Chauhan S, Luorence MJ. The preparation of polyoxyethylene containing non-ionic surfactant vesicles. J Pharm Pharmacol. 1989;41:46.
14. Blazek Welsh AI, Rhodes DG. SEM imaging predicts quality of niosomes from maltodextrin-based proniosomes. Pharm Res. 2001;18:656-61.
15. Kumar AP, Lal J, Jaiswal A, Singh V. Review of niosomes as novel drug delivery system. Int Res J Pharm. 2011; 2(5):61-65.
16. Manosroi A, Wongtrakul P, Manosroi J, Sakai H, Sugawara F, Yuasa M, et al. Characterization of vesicles prepared with various non-ionic surfactants mixed with cholesterol. *Colloids Surf B: Biointerfaces*. 2003;30:129-38.
17. Azmin MN, Florence AT, Handjani-Vila RM, Stuart JF, Vanlerberghe G, Whittaker JS. The effect of non-ionic surfactant vesicle (niosome) entrapment on the absorption and distribution of methotrexate in mice. J Pharm Pharmacol. 1985;37:37-42.
18. Kreuter J. Colloidal drug delivery system of niosome. New York: Marcel Dekker Inc; 1994. p 66-73.
19. Gregoriadis G, Florence AT. In: Liposome technology. Boca Raton, CRC Press, 1993; 2:165.
20. Yoshioka T, Stermberg B, Florence AT. Preparation and properties of vesicles (niosomes) of sobitan monoesters (Span 20, 40, 60, and 80) and a sorbitan triester (Span 85). Int J Pharm. 1994; 105:1-6.
21. Muller RH, Radtke M, Wissing SA. Solid lipid nanoparticles (SLN) and nanostructured lipid carriers (NLC) in cosmetic and dermatological preparations. Adv Drug Deliv. 2002;54: 131-55.
22. Sakthivel M, Kannan K, Manavalan R, Senthamarai R. Formulation and *in vivo* evaluation of niosomes containing oxcarbazepine. J Pharm Sci and Res. 2013;5(1):8-11.
23. Arunothayanun P, Bernard MS, Craig DM, Uchegbu IF, Florence AT. The effect of processing variables on the physical characteristics of non-ionic surfactant vesicles (niosomes) formed from a hexadecyldiglycerol ether. Int J Pharm. 2000;201:7-14.

Review on Niosomes

24. Uchegbu FI, Vyas PS. Non-ionic surfactant based vesicles (niosomes) in drug delivery. *Int J Pharm.* 1998;172:33-70.
25. Muzzalupo R, Tavano L, Lai F, Picci N. Niosomes containing hydroxyl additives as percutaneous penetration enhancers: effect on the transdermal delivery of sulfadiazine sodium salt. *Colloids and Surfaces: B Biointerfaces.* 2014;123:207-212.
26. Tarekegn A, Joseph NM, Palani S, Zacharia A, Ayenew Z. Niosomes in targeted drug delivery: some recent advances. *Int J Phar Sc Res.* 2010; (9):1-8.
27. Stafford S, Ballie AJ, Florence AT. Drug effect on the size of chemically defined nonionic surfactant vesicles. *J Pharm Pharmacol.* 1988;(26):40.
28. Hunter CA, Dolan TF, Coombs GH, Baillie AJ. Vesicular systems (niosomes and liposomes) for delivery of sodium stibogluconate in experimental murine visceral leishmaniasis. *J Pharm Pharmacol.* 1988;40:161-65.
29. Varshosaz J, Pardakhty A, Hajhashemi VI, Najafabadi AR. Development and physical characterization of sorbitan monoester niosomes for insulin oral delivery. *Drug Deliv.* 2003;10:51-62.
30. Brewer JM and Alexander JA. The adjuvant activity of non-ionic surfactant vesicles (niosomes) on the BALB/c humoral response to bovine serum albumin. *Immunology.* 1992; (75):570-75.

Preethi Sudheer and Kaushik Kar

31. Manosroi A, Chankhampan C, Manosroi W, Manosroi J. Transdermal absorption enhancement of papain loaded in elastic niosomes incorporated in gel for scar treatment. *Eu J Pharma Sci.* 2013;48:474-483.
32. Sheena IP, Singh UV, Kamath R, Uma DP, Udupa N. Niosomal withaferin A with better antitumour efficacy. *Indian J Pharm Sci.* 1998;60:45-48.
33. Moser T, Marchand-Arvier M, Labrude P, Vigneron C. Hemoglobin *niosoms*. *Pharm Acta Helv.* 1990;65:82-92.
34. Korkmaz M, Özer AY, Hincal AA. "DTPA niosomes in diagnostic imaging," ed. by Uchegbu I F. Singapore: Harwood Academic Publishers; 2000. p. 227-42.
35. Shahiwala A, Misra A. Studies in topical application of niosomally entrapped nimesulide. *J Pharm Pharm Sci.* 2002;5(3):220-5.
36. Bansal S, Aggarwal G, Chandel P, Harikumar SL. Design and development of cefdinir niosomes for oral delivery. *J Pharm Bioall Sci.* 2013;5(4):318-25.
37. Chaudhari SP, Chatur VM. Development of valproic acid niosomal *in situ* nasal gel formulation for epilepsy. *Ind J Pharma Edu Res.* 2013; 16(3):31-41.